

# Standard Projective Simplicial Kernels and the Second Abelian Cohomology of Topological Groups

H. Sahleh<sup>1</sup> and H.E. Koshkoshi<sup>2</sup>

<sup>1</sup>*Department of Mathematics, Faculty of Mathematical Sciences, University of Guilan  
P. O. Box 1914, Rasht, Iran*

<sup>2</sup>*Department of Mathematics, Faculty of Mathematical Sciences, University of Guilan*

## Abstract

Let  $A$  be an abelian topological  $G$ -module. We give an interpretation for the second cohomology,  $H^2(G, A)$ , of  $G$  with coefficients in  $A$ . As a result we show that if  $P$  is a projective topological group, then  $H^2(P, A) = 0$  for every abelian topological  $P$ -module  $A$ .

**MSC 2010:** Primary 22A05, 20J06, Secondary 18G50.

**Keywords:** Abelian cohomology of topological groups; Markov (Graev) free topological Group; projective topological group; standard projective simplicial kernel.

## 1 Introduction

Let  $G$  be a topological group and  $A$  an abelian topological group. It is said that  $A$  is an abelian topological  $G$ -module, whenever  $G$  acts continuously on  $A$ . For all  $g \in G$  and  $a \in A$  we denote the action of  $g$  on  $a$  by  ${}^ga$ . Suppose that  $A$  is an abelian topological  $G$ -module. Hu [4] showed that if  $G$  is Hausdorff, then there is a one to one correspondence between the second cohomology,  $H^2(G, A)$ , of  $G$  with coefficients in  $A$  and the set,  $Ext_s(G, A)$ , of all equivalence classes of topological extensions with continuous sections. Thus,  $H^2(G, A)$  induces a group structure on  $Ext_s(G, A)$  and consequently, under this group product,  $H^2(G, A)$  is isomorphic to  $Ext_s(G, A)$ . It is known that  $Ext_s(G, A)$  with Baer sum is an abelian group and the Baer sum on  $Ext_s(G, A)$  coincides with the group product induced by  $H^2(G, A)$ . In other words, if  $G$  is Hausdorff then,  $H^2(G, A)$  is isomorphic to  $Ext_s(G, A)$  with Baer sum [1]. Similarly, we define  $Opext_s(G, A)$  and we conclude the same result without Hausdorffness of  $G$ , i.e., there is an isomorphism between  $H^2(G, A)$  and  $Opext_s(G, A)$ . By using the concept of projective topological group we introduce the notion of a standard simplicial kernel of a topological group  $G$ .

in section 2, we prove that every Markov (Graev) topological group is projective and also we show that projectivity in the category of topological groups is equivalent to  $\mathfrak{F}$ -projectivity in the sense of Hall [3]. In addition, we introduce the notion of a standard simplicial kernel of a topological group  $G$ . Finally, we

---

<sup>1</sup>E-mail: [sahleh@guilan.ac.ir](mailto:sahleh@guilan.ac.ir) <sup>2</sup>E-mail: [h.e.koshkoshi@guilan.ac.ir](mailto:h.e.koshkoshi@guilan.ac.ir)

define  $\text{Ope}xt_s(G, A)$  and we conclude that  $H^2(G, A) \cong \text{Ope}xt_s(G, A)$ , with Baer sum.

In section 3, we give a characterization of  $H^2(G, A)$ , when  $A$  is an abelian topological  $G$ -module. As a result, we show that if  $P$  is a projective topological group then  $H^2(P, A) = 0$ , for every abelian topological  $P$ -module.

## 2 Standard Projective Simplicial Kernels.

In this section, we introduce the notion of a standard projective simplicial kernel of a topological group  $G$ .

**Definition 2.1.** A topological group  $P$  is said to be a projective topological group if, for every continuous epimorphism  $\pi : A \rightarrow B$ , where  $\pi$  has a continuous section, and for every continuous homomorphism  $f : P \rightarrow B$ , there exists a continuous homomorphism  $g : P \rightarrow A$  such that the following diagram is commutative

$$\begin{array}{ccc} & P & \\ g \swarrow & \downarrow f & \\ A & \xrightarrow{\pi} B & \longrightarrow 1. \end{array}$$

If  $X$  is a topological space, then the (Markov) free topological group over  $X$  is the pair  $(F_X, \sigma_X : X \rightarrow F_X)$  in which  $F_X$  is a group equipped with the finest group topology such that  $\sigma_X$  is continuous. Such a topology always exists [8], and has the following universal property: every continuous map  $f$  from  $X$  to an arbitrary topological group  $G$  lifts to a unique continuous homomorphism  $\bar{f}$ , i.e.,  $\bar{f} \circ \sigma_X = f$ . Similarly, one can define the Graev free topological group,  $(F_X^*, \sigma_X^*)$ , over a pointed topological space  $(X, e)$ . For information on free (abelian) topological groups see [2, 6, 8]. The following facts about Markov (Graev) free topological groups are well-known:

- (1) *The Markov (Graev) free topological group over a (pointed) topological space  $X$   $((X, e))$  is the free group with the same canonical map over the (pointed) set  $X$   $((X, e))$ ;*
- (2)  *$F_X$   $(F_X^*)$  is Hausdorff if and only if  $X$  is functionally Hausdorff;*
- (3)  *$\sigma_X$   $(\sigma_X^*)$  is a homeomorphic embedding if and only if  $X$  is completely regular;*
- (4)  *$\sigma_X$   $(\sigma_X^*)$  is a closed homeomorphic embedding if and only if  $X$  is Tychonov.*

The elements of free group,  $F_X$ , over a set  $X$  is denoted by  $|x_1|^{\epsilon_1} \dots |x_n|^{\epsilon_n}$ , where  $\epsilon_i = \pm 1$ . This notation is useful whenever  $X$  is a group.

**Lemma 2.2.** *Every Markov (Graev) free topological group is a projective topological group.*

*Proof.* Assume that  $F$  is a Markov free topological group over the space  $X$ , and let  $\pi : A \rightarrow B$  be a continuous epimorphism having a continuous section  $s$  and  $f : F \rightarrow B$  a continuous homomorphism. Then, there is a unique continuous homomorphism  $\tilde{f} : F \rightarrow A$  such that the following diagram is commutative

$$\begin{array}{ccccc} X & \xrightarrow{\sigma_X} & F & \xrightarrow{f} & B \\ \downarrow \sigma_X & & & & \downarrow s \\ F & \xrightarrow{\tilde{f}} & A & & \end{array}$$

Hence,  $sf = \tilde{f}$  on  $X$ . So,  $\pi\tilde{f} = \pi(sf) = (\pi s)f = f$  on  $X$ . Since  $X$  generates the group  $F$ , then it is easy to see that  $\pi\tilde{f} = f$ . Consequently,  $F$  is a projective topological group.

Now, Let  $X$  be a topological space with a fixed point  $e \in X$ . Assume that  $F^*$  is the free topological group in the sense of Graev over  $(X, e)$ , and let  $\pi : A \rightarrow B$  be a continuous epimorphism with continuous section  $s : B \rightarrow A$ . We can assume that  $s$  is a normal section, i.e.,  $s(1) = 1$ , since it is enough to define the new continuous section  $\tilde{s} : B \rightarrow A$  by  $\tilde{s}(b) = s(b)s(1)^{-1}$ . In fact, since  $e$  is the neutral element of  $F^*$ , then  $f(e) = 1$ . Thus,  $sf(e) = 1$ . Hence there exists a unique continuous homomorphism  $\tilde{f} : F^* \rightarrow A$  such that  $sf = \tilde{f}$  on  $X$ . The rest of the proof is the same as the first part.  $\square$

Note that the free functor  $F_{(-)}$  from the category,  $\mathcal{T}$ , of topological spaces to the category,  $\mathcal{TG}$ , of topological groups is left adjoint to the forgetful functor,  $G : \mathcal{TG} \rightarrow \mathcal{T}$ . Let  $\mathfrak{F}$  be the class of all continuous epimorphisms having a continuous section. Thus, by [3, Theorem 2] one can see the following:

*Remark 2.3.* A topological group  $P$  is  $\mathfrak{F}$ -projective if and only if  $P$  is projective.

**Definition 2.4.** Let  $M, F$  and  $G$  be topological groups, and let  $\iota_0, \iota_1 : M \rightarrow F$  and  $\tau : F \rightarrow G$  be continuous homomorphisms. It is said that  $(M, \iota_0, \iota_1)$  is a simplicial kernel of  $\tau$ , whenever  $\tau\iota_0 = \tau\iota_1$  and it has the following universal property:

if  $j_0, j_1 : K \rightarrow F$  are continuous homomorphisms and  $\pi j_0 = \pi j_1$ , then there exists a unique continuous homomorphism  $h : K \rightarrow M$  such that  $j_0 = \iota_0 h$  and  $j_1 = \iota_1 h$ , i.e.,  $h$  commutes the following diagram.

$$\begin{array}{ccccc} K & \xrightarrow{j_0} & F & \xrightarrow{\tau} & G \\ & \searrow j_1 & \parallel & & \\ M & \xrightarrow{\iota_0} & F & \xrightarrow{\tau} & G \\ & \searrow \iota_1 & & & \end{array}$$

**Definition 2.5.** Let  $(M, \iota_0, \iota_1)$  be a simplicial kernel of  $\tau : P \rightarrow G$ . The quadruplet  $(M, \iota_0, \iota_1, \tau)$  is said to be a *projective simplicial kernel of  $G$* , if  $P$  is a projective topological group and  $\tau$  is a continuous (open) epimorphism which has a continuous section  $s : G \rightarrow P$ . If  $P$  is a Markov (Graev) free topological group, then  $(M, \iota_0, \iota_1, \tau)$  is called a *Markov (Graev) simplicial kernel of  $G$* .

**Lemma 2.6.** *If a continuous homomorphism  $f : G \rightarrow H$  has a continuous section, then  $f$  is an open epimorphism.*

*Proof.* The proof is a standard argument.  $\square$

For any topological group  $G$  there exists at least one projective simplicial kernel of  $G$ . Since the identity map  $Id_G : G \rightarrow G$  lifts to a unique homomorphism  $\tau_G : F_G \rightarrow G$ . Thus,  $\tau_G \sigma_G = Id_G$ . The uniqueness property implies that  $\tau_G$  is a map by the following rule:

$$\tau_G : F_G \rightarrow G, |g_1|^{\epsilon_1} \dots |g_n|^{\epsilon_n} \mapsto g_1^{\epsilon_1} \dots g_n^{\epsilon_n} \text{ where } \epsilon_i = \pm 1.$$

The map  $\tau_G$  is called *multiplication map*. Obviously,  $\sigma_G : G \rightarrow F_G, g \mapsto |g|$ , is a continuous section for  $\tau_G$ . By Lemma 2.6,  $\tau_G$  is an open continuous epimorphism. We take  $M_G = \{(x, y) | x, y \in F_G, \tau_G(x) = \tau_G(y)\} \subset F_G \times F_G$ . Hence,  $M_G$  has subspace topology induced by the product topology  $F_G \times F_G$ . Suppose  $\iota_0 = \iota_0^G, \iota_1 = \iota_1^G : M_G \rightarrow F_G$  are the canonical projection maps, where  $\iota_0^G(x, y) = x$  and  $\iota_1^G(x, y) = y$ . It is easy to see that  $(M_G, \iota_0^G, \iota_1^G)$  is a simplicial kernel of  $\tau_G$ . Hence,  $(M_G, \iota_0^G, \iota_1^G, \tau_G)$  is a projective simplicial kernel of  $G$  and we call it the *standard Markov simplicial kernel of  $G$* . By a similar way we can define the *standard Graev (projective) simplicial kernel of  $G$* .

Suppose that  $(M, \iota_0, \iota_1, \tau)$  is a projective simplicial kernel of  $G$ . We sometimes denoted it by the following

$$M \begin{array}{c} \xrightarrow{\iota_0} \\ \xrightarrow{\iota_1} \end{array} P \xrightarrow{\tau} G \longrightarrow 1. \quad (*)$$

If  $(M_G, \iota_0, \iota_1, \tau_G)$  is a standard projective simplicial kernel of  $G$ , then put  $\Delta_P = \{(x, x) | x \in P\} \subset M$ . Now, let  $A$  be an abelian topological  $G$ -module. Obviously,  $A$  is a topological  $P$ -module via  $\tau$  and a topological  $M$ -module via  $\tau\iota_0$  (or  $\tau\iota_1$ ). Define  $Z^1(M, A) = \{\alpha | \alpha \in Der_c(M, A), \alpha(\Delta_P) = 0\}$ . It is clear that  $Z^1(M, A)$  is a subgroup of abelian group  $Der_c(M, A)$ .

Consider

$$\begin{aligned} \eta : Der_c(P, A) &\rightarrow Z^1(M, A) \\ \alpha &\mapsto \alpha\iota_0\alpha\iota_1^{-1} \end{aligned}$$

Obviously,  $\eta$  is well-defined and since  $A$  is an abelian group, then

$$\begin{aligned} \eta(\alpha.\beta) &= (\alpha.\beta\iota_0)(\alpha.\beta\iota_1^{-1}) = (\alpha\iota_0.\beta\iota_0)(\alpha\iota_1^{-1}.\beta\iota_1^{-1}) \\ &= (\alpha\iota_0\alpha\iota_1^{-1})(\beta\iota_0\beta\iota_1^{-1}) = \eta(\alpha)\eta(\beta). \end{aligned}$$

Therefore,  $\eta$  is a homomorphism. Clearly,  $\eta$  induces a congruence equivalence relation  $\sim$  in  $Z^1(M, A)$ . In other words, for  $\alpha, \beta \in Z^1(M, A)$ , the equivalence relation  $\sim$  is defined as follows:

$$\alpha \sim \beta \text{ whenever there is } \gamma \in Der_c(P, A) \text{ and } \beta = \gamma\iota_0\gamma\iota_1^{-1}\alpha.$$

**Definition 2.7.** It is said that a short exact sequence

$$e : 1 \rightarrow A \xrightarrow{\chi} E \xrightarrow{\pi} G \rightarrow 1$$

is a proper extension of  $G$  by  $A$  whenever  $\chi$  is a homeomorphic embedding and  $\pi$  is an open continuous homomorphism. By a section for  $e$  we mean a continuous map  $s : G \rightarrow E$  such that  $\pi s = Id_G$ .

Let  $e$  be a proper extension of  $G$  by  $A$  which has a section  $s : G \rightarrow E$ . In addition, assume that  $A$  is abelian. Then the proper extension  $e$  gives rise to a topological  $G$ -module structure on  $A$ , which is well-defined by

$${}^g a = \chi^{-1}(s(g)\chi(a)s(g)^{-1}) , \text{ where } g \in G , a \in A.$$

One can easily see that for each  $g \in G$  and  $a \in A$ , the element  ${}^g a$  dose not depend on the choice of continuous sections.

Now, let  $A$  be an abelian topological  $G$ -module. We denote by  $opext_s(G, A)$  the set of all proper extensions of  $G$  by  $A$  having continuous sections and corresponding to the given way in which  $G$  acts on  $A$ . We define an equivalence relation  $\equiv$  on  $opext_s(G, A)$  as follows:

Let  $e_i : 1 \rightarrow A \xrightarrow{\chi_i} E \xrightarrow{\pi_i} G \rightarrow 1 \in opext_s(G, A)$  for  $i = 0, 1$ .  $e_0 \equiv e_1$  whenever there is a continuous (open) homomorphism  $\theta : E_0 \rightarrow E_1$  so that the following diagram is commutative.

$$\begin{array}{ccccccccc} 1 & \longrightarrow & A & \xrightarrow{\chi_0} & E_0 & \xrightarrow{\pi_0} & G & \longrightarrow & 1 \\ & & \parallel & & \downarrow \theta & & \parallel & & \\ 1 & \longrightarrow & A & \xrightarrow{\chi_1} & E_1 & \xrightarrow{\pi_1} & G & \longrightarrow & 1 \end{array}$$

We denote the set of all equivalence classes by  $Opext_s(G, A)$ . Note that if  $G$  is Hausdorff, then  $Opext_s(G, A) = Ext_s(G, A)$ . One can verify by a similar argument as in [1, 4] that we have the following.

**Theorem 2.8.** *Let  $A$  be an abelian topological  $G$ -module, then  $H^2(G, A)$  in the sense of Hu, is isomorphic to  $Opext_s(G, A)$  with Baer sum.*

### 3 the Main Theorem

In this section we will prove the main theorem. Sometimes we denote the inclusion maps and epics by the arrows  $\hookrightarrow$  and  $\twoheadrightarrow$ , respectively. Let  $A$  be a topological  $G$ -module. Thus, under the action  $(g, a) \mapsto {}^g a$  we may consider the topological semidirect product  $A \rtimes_\phi G$  (see [7, Section 6]).

**Theorem 3.1.** *Let  $A$  be an abelian topological  $G$ -module and let  $(M, \iota_0, \iota_1, \tau)$  be a standard projective simplicial kernel of  $G$ . Then,  $H^2(G, A)$  is canonically isomorphic to  $Z^1(M, A)/\sim$ .*

*Proof.* By Theorem 2.8, it is enough to show that  $\text{Ope}xt_s(G, A)$  is isomorphic to  $Z^1(M, A)/\sim$ .

Let  $\alpha \in Z^1(M, A)$ . Since  $A$  is a  $G$ -module, then  $A$  is a  $P$ -module via  $\tau$ . Thus, we may consider topological semidirect product  $A \rtimes P$ . Now, we define a relation  $\sim$  in the semidirect product  $A \rtimes P$ , as follows

$$(a, x) \sim (b, y) \text{ if and only if } \tau(x) = \tau(y) \text{ and } a\alpha(x, y) = b.$$

Obviously,  $\sim$  is reflexive. Let  $x, y$  and  $z$  be arbitrary elements of  $P$  such that  $\tau(x) = \tau(y) = \tau(z)$ . Then

$$\alpha(x, y)\alpha(y, z) = \alpha(x, z) \quad (3.1)$$

since,

$$\begin{aligned} \alpha(x, z) &= \alpha((x, y)(1, y^{-1}z)) = \alpha(x, y) \cdot^y \alpha(1, y^{-1}z) = \alpha(x, y) \cdot^y \alpha(y^{-1}y, y^{-1}z) \\ &= \alpha(x, y) \cdot^y \alpha((y^{-1}, y^{-1})(y, z)) = \alpha(x, y) \cdot^y (\alpha(y^{-1}, y^{-1}) \cdot^{y^{-1}} \alpha(y, z)) \\ &= \alpha(x, y) \cdot^y (1 \cdot^{y^{-1}} \alpha(y, z)) = \alpha(x, y)\alpha(y, z). \end{aligned}$$

Now if  $(a, x) \sim (b, y)$  and  $(b, y) \sim (c, z)$ , then

$$\tau(x) = \tau(y) = \tau(z), a\alpha(x, y) = b \text{ and } b\alpha(y, z) = c.$$

Hence, by (3.1),  $a\alpha(x, z) = a\alpha(x, y)\alpha(y, z) = b\alpha(y, z) = c$ . This implies that  $\sim$  is transitive. By (3.1), we have:

$$\alpha^{-1}(x, y) = \alpha(y, x). \quad (3.2)$$

If  $(a, x) \sim (b, y)$ , then  $\tau(x) = \tau(y)$  and  $a\alpha(x, y) = b$ . Thus,  $a = b\alpha^{-1}(x, y) = b\alpha(y, x)$ . Consequently,  $\sim$  is an equivalence relation in  $A \rtimes P$ .

Now, Define  $N_\alpha = \{(a, x) | (a, x) \in A \rtimes P, (a, x) \sim (1, 1)\}$ . Then, by the definition of  $\sim$  and the identity (3.2), we have

$$\begin{aligned} N_\alpha &= \{(a, x) | (a, x) \in A \rtimes P, \tau(x) = 1, a\alpha(x, 1) = 1\} \\ &= \{(a, x) | (a, x) \in A \rtimes P, \tau(x) = 1, a = \alpha(1, x)\}. \end{aligned}$$

Suppose that  $(a, x), (b, x) \in N_\alpha$ . Since  $\tau(x) = 1$ , then  $(a, x)(b, x) = (a^x b, xy) = (ab, xy)$ . On the other hand,  $\alpha(1, xy) = \alpha(1, x)\alpha(1, y) = ab$ . Thus,  $(a, x)(b, x) \in N_\alpha$ . Also  $(a, x)^{-1} = (x^{-1}a^{-1}, x^{-1}) = (a^{-1}, x^{-1})$  and  $\alpha(1, x^{-1}) = \alpha(xx^{-1}, x^{-1}) = \alpha(x, 1)^x \alpha(x^{-1}, x^{-1}) = \alpha(x, 1) = \alpha^{-1}(1, x) = a^{-1}$ , hence  $(a, x)^{-1} \in N_\alpha$ . Therefore  $N_\alpha$  is a subgroup of  $A \rtimes P$ .

Denote by  $E_\alpha$  the quotient space  $(A \rtimes P)/\sim$ .

We show that  $\sim$  is a congruence (i.e.,  $\sim$  is compatible with the group product) and therefore  $E_\alpha$  is a group.

Let  $(a, x) \sim (a', x')$  and  $(b, y) \sim (b', y')$ . Then  $\tau(x) = \tau(x')$ ,  $\tau(y) = \tau(y')$ ,  $a\alpha(x, x') = a'$  and  $b\alpha(y, y') = b'$ .

We have  $(a, x)(b, y) = (a^x b, xy)$ ,  $(a', x')(b', y') = (a'^{x'} b', x'y')$ . Since  $A$  is abelian, then

$$a^x b \alpha(xy, x'y') = a^x b \alpha(x, x')^x \alpha(y, y') = a \alpha(x, x')^x (b \alpha(y, y')) = a'^x b' = a'^{x'} b'.$$

Hence,  $(a, x)(b, y) \sim (a', x')(b', y')$ . Thus,  $E_\alpha$  is a group. It is clear that  $E_\alpha = (A \rtimes P)/N_\alpha$ . This implies that  $N_\alpha$  is a normal subgroup of  $A \rtimes P$ . Therefore  $E_\alpha$  is a topological group.

There is a diagram as follows:

$$\begin{array}{ccccccc} M & \xrightarrow{\iota_0} & P & \xrightarrow{\tau} & G & \longrightarrow & 1 \\ & \searrow \iota_1 & \downarrow \beta & & \parallel & & \\ A & \xrightarrow{\sigma} & E_\alpha & \xrightarrow{\psi} & G & \longrightarrow & 1 \end{array} \quad (**)$$

where  $\sigma(a) = (a, 1)N_\alpha$ ,  $\psi(a, x)N_\alpha = \tau(x)$  and  $\beta(x) = (1, x)N_\alpha$ .

It is obvious that  $\sigma$  and  $\beta$  are continuous homomorphisms. Consider the continuous map  $\phi : A \rtimes P \rightarrow P$ ,  $(a, x) \mapsto \tau(x)$ .

We know that  $\tau$  has a continuous section  $s : G \rightarrow P$ . Hence,  $\beta \circ s$  is a continuous section for  $\psi$ , since  $\psi \beta \circ s = (\psi \beta) \circ s = \tau \circ s = Id_G$ . Thus, by Lemma 1.6,  $\phi$  is an open epimorphism.

If  $\sigma(a) \in N_\alpha$ , then  $(a, 1) \in N_\alpha$ . Thus, by definition of  $N_\alpha$ , we get  $a = \alpha(1, 1) = 1$ . Therefore  $\sigma$  is one to one.

$\psi \circ \sigma(a) = \psi((a, 1)N_\alpha) = \tau(1) = 1$ , thus  $Im\sigma \subset Ker\psi$ . On the other hand, if  $\psi(a, x)N_\alpha = 1$ , then  $\tau(x) = 1$  and  $\sigma(a\alpha(x, 1)) = (a, x)N_\alpha$ , i.e.,  $Ker\psi \subset Im\sigma$ . Therefore,  $Im\sigma = Ker\psi$ .

Now, we prove that  $\sigma$  is a homeomorphic embedding.

Define the map  $\chi : A \rtimes P \rightarrow A$  via  $\chi(a, x) = a\alpha(x, s\tau(x)s(1)^{-1})$ . Obviously,  $\chi$  is continuous. It is clear that  $(A \times \{1\})N_\alpha$  is a topological subgroup of  $A \rtimes P$  and  $(A \times \{1\})N_\alpha = \{(a, x) | a \in A, x \in Ker\tau\}$ . Take  $\hat{\chi} = \chi|_{(A \times \{1\})N_\alpha}$ . Thus,  $\hat{\chi}$  is continuous and  $\hat{\chi}(a, x) = a\alpha(x, 1)$ . Note that  $\hat{\chi}$  is a homomorphism, because

$$\begin{aligned} \hat{\chi}((a, x)(b, y)) &= \hat{\chi}(ab, xy) = ab\alpha(xy, 1) = ab\alpha(x, 1)\alpha(xy, x) \\ &= ab\alpha(x, 1)\alpha(x^{-1}xy, 1) = ab\alpha(x, 1)\alpha(y, 1) = \hat{\chi}(a, x)\hat{\chi}(b, y). \end{aligned}$$

If  $(a, x) \in N_\alpha$ , then  $\hat{\chi}(a, x) = a\alpha(x, 1) = \alpha(1, x)\alpha(x, 1) = \alpha(1, 1) = 1$ . Hence  $\hat{\chi}(N_\alpha) = 1$ . So,  $\hat{\chi}$  induces the continuous homomorphism

$$\xi : ((A \times \{1\})N_\alpha)/N_\alpha \rightarrow A, \xi((a, x)N_\alpha) = \hat{\chi}(a, x).$$

It is clear that  $Im\sigma = ((A \times \{1\})N_\alpha)/N_\alpha$ . We have  $\xi \circ \sigma(a) = a$  and

$$\begin{aligned} \sigma \circ \xi((a, x)N_\alpha) &= \sigma(a\alpha(x, 1)) = (a\alpha(x, 1), 1)N_\alpha = (a, x)(\alpha(x, 1), x^{-1})N_\alpha \\ &= (a, x)(\alpha(1, x^{-1}), x^{-1})N_\alpha = (a, x)N_\alpha. \end{aligned}$$

Thus,  $\sigma|^{Im\sigma} : A \rightarrow Im\sigma$ ,  $\sigma|^{Im\sigma}(a) = (a, 1)N_\alpha$ , is a topological isomorphism. Therefore,  $\sigma$  is a homeomorphic embedding.

Denote by  $e_\alpha$  the extension  $0 \rightarrow A \xrightarrow{\sigma} E_\alpha \xrightarrow{\psi} G \rightarrow 1$ . We have  $[e_\alpha] \in \text{Ope}xt_s(G, A)$ . Since  $s : G \rightarrow P$  is a continuous section for  $\tau$ , then  $\beta \circ s$  is a continuous section for  $\psi$  and

$$\begin{aligned} \beta \circ s(g)\sigma(a)\beta \circ s(g)^{-1} &= (1, s(g))N_\alpha(a, 1)N_\alpha(1, s(g)^{-1})N_\alpha \\ &= (1, s(g))(a, 1)(1, s(g)^{-1})N_\alpha = ({}^{s(g)}a, s(g))(1, s(g)^{-1})N_\alpha \\ &= ({}^{s(g)}a, 1)N_\alpha = ({}^ga, 1)N_\alpha = \sigma({}^ga). \end{aligned}$$

In addition,  $\sigma$  is one to one, thus the extension  $e_\alpha$  induces the given action of  $G$  on  $A$ . Hence  $[e_\alpha] \in \text{Ope}xt_s(G, A)$ .

Consider the map  $\zeta : Z^1(M, A) \rightarrow \text{Ope}xt_s(G, A)$  via  $\alpha \mapsto [e_\alpha]$ . Suppose that  $\alpha, \bar{\alpha} \in Z^1(M, A)$  and  $\bar{\alpha} = \eta(\gamma)\alpha$ , for some  $\gamma \in \text{Der}_c(P, A)$ . Define the map  $\varepsilon : A \rtimes P \rightarrow A \rtimes P$  by  $(a, x) \mapsto (a\gamma^{-1}(x), x)$ . Obviously,  $\varepsilon$  is continuous, and

$$\begin{aligned} \varepsilon((a, x)(b, y)) &= \varepsilon(a^xb, xy) = (a^xb\gamma^{-1}(xy), xy) = (a^xb^x\gamma^{-1}(y)\gamma^{-1}(x), xy) \\ &= (a\gamma^{-1}(x)^x(b\gamma^{-1}(y)), xy) = (a\gamma^{-1}(x), x)(b\gamma^{-1}(y), y); \end{aligned}$$

that is,  $\varepsilon$  is a homomorphism.

If  $(a, x) \in N_\alpha$ , then

$$\begin{aligned} \bar{\alpha}(1, x) &= (\gamma\iota_0 \cdot \gamma\iota_1^{-1} \cdot \alpha)(1, x) = \gamma\iota_0(1, x)\gamma\iota_1^{-1}(1, x)\alpha(1, x) = \gamma^{-1}(x)\alpha(1, x) \\ &= a\gamma^{-1}(x). \end{aligned}$$

Hence,  $(a\gamma^{-1}(x), x) \in N_{\bar{\alpha}}$ , i.e.,  $\varepsilon(N_\alpha) \subset N_{\bar{\alpha}}$ . Thus,  $\varepsilon$  induces the continuous homomorphism  $\hat{\varepsilon} : E_\alpha \rightarrow E_{\bar{\alpha}}$ ,  $(a, x)N_\alpha \mapsto (a\gamma^{-1}(x), x)N_{\bar{\alpha}}$ .

Suppose that the extensions  $e_\alpha : 0 \rightarrow A \xrightarrow{\sigma} E_\alpha \xrightarrow{\psi} G \rightarrow 1$  and  $e_{\bar{\alpha}} : 0 \rightarrow A \xrightarrow{\bar{\sigma}} E_{\bar{\alpha}} \xrightarrow{\bar{\psi}} G \rightarrow 1$  are corresponding to  $\alpha$  and  $\bar{\alpha}$ , respectively. We show that the following diagram commutes.

$$\begin{array}{ccccccc} e_\alpha : 0 & \longrightarrow & A & \xrightarrow{\sigma} & E_\alpha & \xrightarrow{\psi} & G \longrightarrow 1 \\ & & \parallel & & \downarrow \hat{\varepsilon} & & \parallel \\ e_{\bar{\alpha}} : 0 & \longrightarrow & A & \xrightarrow{\bar{\sigma}} & E_{\bar{\alpha}} & \xrightarrow{\bar{\psi}} & G \longrightarrow 1 \end{array}$$

Because,  $\bar{\psi}\hat{\varepsilon}(a, x)N_\alpha = \bar{\psi}(a\gamma^{-1}(x), x)N_{\bar{\alpha}} = \tau(x) = \psi(a, x)N_\alpha$ , and  $\hat{\varepsilon}\sigma(a) = \hat{\varepsilon}(a, 1)N_\alpha = (a\gamma^{-1}(1), 1)N_{\bar{\alpha}} = (a, 1)N_{\bar{\alpha}} = \bar{\sigma}(a)$ . i.e.,  $[e_\alpha] = [e_{\bar{\alpha}}]$ . Thus,  $\zeta$  induces the map

$$\begin{aligned} \Theta : \text{Coker}\eta &\rightarrow \text{Ope}xt_s(G, A) \\ [\alpha] &\mapsto [e_\alpha]. \end{aligned}$$

We will prove that  $\Theta$  is a bijective map.

Let  $e : 0 \longrightarrow A \xrightarrow{\sigma} C \xrightarrow{\psi} G \longrightarrow 1$  be an extension with a continuous section  $s : G \rightarrow A$  and corresponding to the given way in which  $G$  acts

on  $A$ . Since  $P$  is a projective topological group, then there is a continuous homomorphism  $\beta$  so that

$$\begin{array}{ccccc} P & \xrightarrow{\tau} & G & \longrightarrow & 1 \\ \downarrow \beta & & \parallel & & \\ C & \xrightarrow{\psi} & G & \longrightarrow & 1 \end{array}$$

commutes. Now, define  $\alpha_\beta : M \rightarrow A$  by  $\alpha_\beta(m) = (\beta\iota_0\beta\iota_1^{-1})(m)$ . Note that  $\alpha_\beta$  is well-defined, because  $\psi(\beta\iota_0\beta\iota_1^{-1}) = \psi\beta\iota_0\psi\beta\iota_1^{-1} = \tau\iota_0\tau\iota_1^{-1} = \mathbf{1}$ . Thus,  $\text{Im}\alpha_\beta \subset \text{Ker}\psi = \text{Im}\sigma = A$ . For simplicity, we denote  $\alpha_\beta$  by  $\beta\iota_0\beta\iota_1^{-1}$ . It is clear that  $\alpha_\beta$  is continuous and in addition,  $\alpha_\beta$  is a crossed homomorphism, since

$$\begin{aligned} \alpha_\beta((x_1, y_1)(x_2, y_2)) &= \alpha_\beta(x_1x_2, y_1y_2) = \beta(x_1x_2)\beta^{-1}(y_1y_2) \\ &= \beta(x_1)\beta(x_2)\beta^{-1}(y_2)\beta^{-1}(y_1) = \beta(x_1)\alpha_\beta(x_2, y_2)\beta^{-1}(y_1) \\ &= \alpha_\beta(x_1, y_1)\beta(y_1)\alpha_\beta(x_2, y_2)\beta^{-1}(y_1) = \alpha_\beta(x_1, x_2)^{\beta(y_1)}\alpha_\beta(x_2, y_2) \\ &= \alpha_\beta(x_1, y_1)^{\psi\beta(y_1)}\alpha_\beta(x_2, y_2) = \alpha_\beta(x_1, y_1)^{\tau(y_1)}\alpha_\beta(x_2, y_2) \\ &= \alpha_\beta(x_1, y_1)^{(x_1, y_1)}\alpha_\beta(x_2, y_2), \end{aligned}$$

and  $\alpha_\beta(x, x) = \beta(x)\beta^{-1}(x) = 1$ . i.e.,  $\alpha_\beta(\Delta_P) = 1$ , hence  $\alpha_\beta \in Z^1(M, A)$ . Consequently,  $\beta$  and  $\alpha_\beta$  make the following diagram commutative.

$$\begin{array}{ccccccc} M & \xrightarrow{\iota_0} & P & \xrightarrow{\tau} & G & \longrightarrow & 1 \\ \downarrow \alpha_\beta & \swarrow \iota_1 & \downarrow \beta & & \parallel & & \\ 0 \longrightarrow & A & \xrightarrow{\sigma} & C & \xrightarrow{\psi} & G & \longrightarrow 1 \end{array}$$

Now, we will show that  $[\alpha_\beta]$  is independent of the choice of  $\beta : P \rightarrow C$ . Consider the commutative diagram.

$$\begin{array}{ccccccc} M & \xrightarrow{\iota_0} & P & \xrightarrow{\tau} & G & \longrightarrow & 1 \\ \downarrow \alpha_{\bar{\beta}} & \swarrow \iota_1 & \downarrow \bar{\beta} & & \parallel & & \\ 0 \longrightarrow & A & \xrightarrow{\sigma} & C & \xrightarrow{\psi} & G & \longrightarrow 1 \end{array}$$

Define  $\gamma(x) = (\bar{\beta}\beta^{-1})(x)$ , for all  $x \in P$ . Since  $\psi\gamma(x) = \psi\bar{\beta}(x)\psi\beta^{-1}(x) = \tau(x)\tau(x)^{-1} = 1$ , then  $\gamma(x) \in \text{Ker}\psi = A$ . Hence, we can define the continuous map  $\gamma : P \rightarrow A$ ,  $x \mapsto \gamma(x)$ . Now  $\gamma$  is a crossed homomorphism, since

$$\begin{aligned} \gamma(xy) &= (\bar{\beta}\beta^{-1})(xy) = \bar{\beta}(xy)\beta^{-1}(xy) = \bar{\beta}(x)\bar{\beta}(y)\beta^{-1}(y)\beta^{-1}(x) \\ &= \bar{\beta}(x)\gamma(y)\beta^{-1}(x) = \gamma(x)\beta(x)\gamma(y)\beta^{-1}(x) = \gamma(x)^{\beta(x)}\gamma(y) \\ &= \gamma(x)^{\psi\beta(x)}\gamma(y) = \gamma(x)^x\gamma(y). \end{aligned}$$

We have  $\bar{\beta} = \gamma\beta$ . Thus,

$$\alpha_{\bar{\beta}} = \bar{\beta}\iota_0\bar{\beta}\iota_1^{-1} = \gamma\beta\iota_0\gamma\beta\iota_0^{-1} = \gamma\iota_0\beta\iota_0\beta\iota_1^{-1}\gamma\iota_1^{-1} = \gamma\iota_0(\beta\iota_0\beta\iota_1^{-1})\gamma\iota_1^{-1} = \gamma\iota_0\alpha_{\beta}\gamma\iota_1^{-1} = \alpha_{\beta}(\gamma\iota_0\gamma\iota_1^{-1}).$$

i.e.,  $\alpha_{\bar{\beta}} \sim \alpha_{\beta}$ .

Now, we will show that  $\Theta([\alpha_{\beta}]) = [e]$ . It is sufficient to prove that  $e_{\alpha_{\beta}} \sim e$ . Define the map  $\nu : A \rtimes P \rightarrow C$  via  $(a, x) \mapsto a\beta(x)$ . Obviously,  $\nu$  is continuous, and since

$$\begin{aligned} \nu((a, x)(b, y)) &= \nu(a^x b, xy) = a^x b\beta(xy) = a\beta(x)^{\beta(x)^{-1}}(xb)\beta(y) = \\ &= \nu(a, x)^{\psi\beta(x)^{-1}}(xb)\beta(y) = \nu(a, x)^{x^{-1}}(xb)\beta(y) = \nu(a, x)\nu(b, y), \end{aligned}$$

then  $\nu$  is a homomorphism. On the other hand, if  $(a, x) \in N_{\alpha_{\beta}}$ , then

$$\nu(a, x) = a\beta(x) = \alpha_{\beta}(1, x)\beta(x) = \beta^{-1}(x)\beta(x) = 1.$$

Thus,  $\nu : A \rtimes P \rightarrow C$  induces the continuous homomorphism  $\hat{\nu} : E_{\alpha_{\beta}} \rightarrow C$ ,  $(a, x)N_{\alpha_{\beta}} \mapsto a\beta(x)$ . Note that  $\hat{\nu}$  commutes the following diagram

$$\begin{array}{ccccccc} e_{\alpha_{\beta}} : 0 & \longrightarrow & A & \xrightarrow{\bar{\sigma}} & E_{\alpha_{\beta}} & \xrightarrow{\bar{\psi}} & G \longrightarrow 1 \\ & & \parallel & & \downarrow \hat{\nu} & & \parallel \\ e : 0 & \longrightarrow & A & \xrightarrow{\sigma} & C & \xrightarrow{\psi} & G \longrightarrow 1. \end{array}$$

Since  $\hat{\nu}\bar{\sigma}(a) = \hat{\nu}((a, 1)N_{\alpha_{\beta}}) = a\beta(1) = a = \sigma(a)$ , and  $\psi\hat{\nu}((a, x)N_{\alpha_{\beta}}) = \psi(a\beta(x)) = \psi\beta(x) = \tau(x) = \psi((a, x)N_{\alpha_{\beta}})$ . Therefore,  $e_{\alpha_{\beta}} \sim e$ . i.e.,  $\Theta$  is onto.

We show that  $\Theta$  is one to one.

Let  $e_i : 0 \longrightarrow A \xrightarrow{\sigma_i} C_i \xrightarrow{\psi_i} G \longrightarrow 1$ ,  $i = 0, 1$ , be the extensions with continuous sections and corresponding to the given way in which  $G$  acts on  $A$ . Suppose that  $e_1 \sim e_2$  are equivalent, i.e., there is continuous homomorphism  $\omega : C_1 \rightarrow C_2$  such that  $\omega(a) = a, \forall a \in A$ , and  $\psi_2\omega = \psi_1$ . There is a continuous homomorphism  $\beta : P \rightarrow C_1$  such that  $\psi_1\beta = \tau$ . Take  $\bar{\beta} = \omega \circ \beta$ . Consider the following diagram.

$$\begin{array}{ccccccc} & & M & \xrightarrow{\iota_0} & P & \xrightarrow{\tau} & G \longrightarrow 1 \\ & & \downarrow \alpha_{\beta} & & \downarrow \beta & & \parallel \\ \alpha_{\bar{\beta}} \nearrow & 0 \longrightarrow & A & \xrightarrow{\sigma_1} & C_1 & \xrightarrow{\psi_1} & G \longrightarrow 1 \\ & & \downarrow \alpha_{\bar{\beta}} & & \downarrow \bar{\beta} & & \parallel \\ & & A & \xrightarrow{\sigma_2} & C_2 & \xrightarrow{\psi_2} & G \longrightarrow 1 \end{array}$$

Since the back, the bottom, the front squares, and the middle triangle are commutative, so is the left triangle, i.e.,  $\alpha_{\bar{\beta}} = \alpha_{\beta}$ . This shows that  $\Theta$  is one to one. Consequently,  $\Theta$  is bijective.

Finally, we prove that  $\Theta$  is a homomorphism. Suppose that  $\Theta([\alpha_i]) = [e_i]$ ,  $i = 0, 1$ . In another words, the following diagram is commutative.

$$\begin{array}{ccccccc}
 & M & \xrightarrow{\iota_0} & P & \xrightarrow{\tau} & G & \longrightarrow 1 \\
 & \downarrow \alpha_i & \searrow \iota_1 & \downarrow \beta_i & & \parallel & \\
 e_i : 0 & \longrightarrow & A & \xrightarrow{\sigma_i} & E_i & \xrightarrow{\psi_i} & G \longrightarrow 1
 \end{array}$$

for  $i = 0, 1$ . Hence, we have the commutative diagram

$$\begin{array}{ccccccc}
 & M & \xrightarrow{\iota_0} & P & \xrightarrow{\tau} & G & \longrightarrow 1 \\
 & \downarrow \alpha_1 \times \alpha_2 & \searrow \iota_1 & \downarrow \beta_1 \times \beta_2 & & \Delta_G \downarrow & \\
 e_1 \times e_2 : 0 & \longrightarrow & A \times A & \xrightarrow{\sigma_1 \times \sigma_2} & E_1 \times E_2 & \xrightarrow{\psi_1 \times \psi_2} & G \times G \longrightarrow 1
 \end{array}$$

where  $\Delta_G(g) = (g, g)$ ,  $\beta_1 \times \beta_2(p) = (\beta_1(p), \beta_2(p))$ ,  $\alpha_1 \times \alpha_2(m) = (\alpha_1(m), \alpha_2(m))$ ,  $\sigma_1 \times \sigma_2(a, b) = (\sigma_1(a), \sigma_2(b))$  and  $\psi_1 \times \psi_2(e_1, e_2) = (\psi_1(e_1), \psi_2(e_2))$ . Then by definition of pushout and pullback extensions [Al], we get the following diagram

$$\begin{array}{ccccccc}
 & M & \xrightarrow{\iota_0} & P & \xrightarrow{\tau} & G & \\
 & \downarrow \alpha_1 \cdot \alpha_2 & \searrow \iota_1 & \downarrow \beta_1 \times \beta_2 & & \parallel & \\
 e_4 : & A & \xrightarrow{\quad} & E_4 & \xrightarrow{\quad} & G & \\
 & \downarrow \nabla_A & \searrow & \downarrow & & \Delta_G \downarrow & \\
 & A \times A & \xrightarrow{\sigma_1 \times \sigma_2} & E_1 \times E_2 & \xrightarrow{\psi_1 \times \psi_2} & G \times G & \\
 & \downarrow & \searrow & \downarrow & & \parallel & \\
 e_3 : & A & \xrightarrow{\quad} & E_3 & \xrightarrow{\quad} & G \times G & 
 \end{array} \quad (***)$$

in which, extension  $e_3 = \nabla_A(e_1 \times e_2) : 0 \longrightarrow A \longrightarrow E_3 \longrightarrow G \times G \longrightarrow 1$  is the pushout extension of  $e_1 \times e_2$ , and extension  $e_4 = (\nabla_A(e_1 \times e_2))\Delta_G : 0 \longrightarrow A \longrightarrow E_4 \longrightarrow G \longrightarrow 1$  is the pullback extension of  $\nabla_A(e_1 \times e_2)$ . Also,  $\nabla_A$  is the codiagonal map, i.e.,  $\nabla_A(a, b) = ab$ , and  $\alpha_1 \cdot \alpha_2(m) = \alpha_1(m)\alpha_2(m)$ . Obviously, the left and the right squares are commutative. Since  $P$  is a projective topological group, then the existence of  $\gamma : P \rightarrow A$  is guaranteed and commutes the right-up square. Therefore in the right cube of diagram (\*\*\*), all the up, down, front, back and right squares are commutative. So  $\gamma$  commutes the middle square. On the other hand, in the left cube of diagram (\*\*\*), all the right, left, back, front and down squares are commutative. Thus,  $\gamma$  commutes the left-up square of diagram (\*\*\*). Therefore, the whole top of diagram (\*\*\*) is commutative. This means that  $\Theta([\alpha_1 \cdot \alpha_2]) = [e_4]$ . On the other hand by

definition of Baer sum in the  $Opext_s(G, A)$ ,  $[\nabla_A(e_1 \times_2) \Delta_G] = [e_1] + [e_2]$ . Consequently,  $\Theta(\alpha_1 \cdot \alpha_2) = [e_1] + [e_2] = \Theta(\alpha_1) + \Theta(\alpha_2)$ , i.e.,  $\Theta$  is an isomorphism and thereby  $\Theta$  is an isomorphism. The proof is completed.  $\square$

Note that Theorem 3.1 is similar to [5, Proposition 6] in a topological context.

**Corollary 3.2.** *Let  $P$  be a projective topological group, then  $H^2(P, A) = 0$ , for every abelian topological  $P$ -module  $A$ . In particular, for every Markov (Graev) free topological groups  $F$ ,  $H^2(F, A) = 0$ , for every abelian topological  $F$ -module  $A$ .*

*Proof.* It is clear that  $(\Delta_P, \iota_0, \iota_G, Id_P)$  is a standard projective simplicial kernel of  $P$ . Thus, every continuous crossed homomorphism  $\alpha : \Delta_P \rightarrow A$  is the zero map. Consequently,  $H^2(P, A) = 0$ .  $\square$

Thus, we have discovered a general result of [4, (5.6)].

## References

- [1] R.C. Alperin and H. Sahleh, Hopf's formula and the Schur multiplier for topological groups, Kyungpook. Math. Journal, 31(1) (1991), 35-71.
- [2] M. I. Graev, Free topological groups, (Russian) Izv. Akad. Nauk SSSR, Ser. Mat. 12 (1948), 279-324.
- [3] C. E. Hall,  $\mathfrak{F}$ -Projective Objects, Proc. Amer. Math. Soc. 26, (1970), 193-195.
- [4] S.T. Hu, Cohomology theory in topological groups, Michigan Math. J., 1(1) (1952), 11-59.
- [5] H. Inassaridze, Non-abelian cohomology of groups, Georgian Math. J. 4, No 4 (1997), 313-332.
- [6] A. A. Markov, On free topological groups. (Russian) Izv. Akad. Nauk SSSR, Ser. Mat. 9 (1945), 3-64.
- [7] H. Sahleh, H.E. Koshkoshi, First Non-Abelian Cohomology Of Topological Groups. Gen. Math. Notes, 2012, 12(1), 20-34.
- [8] B.V.S. Thomas, Free topological groups, General Topology and its Appl., 1 (1971), pp. 5172.